


Measuring the Acceleration of the Cosmic Expansion Using Supernovae

Saul Perlmutter


*University of California, Berkeley
Lawrence Berkeley National Laboratory*

Nobel Lecture
Stockholm
December 2011


A deep space photograph showing a vast field of stars and galaxies against a black background. The stars vary in size, color (white, yellow, blue), and brightness. Some galaxies are visible as faint, irregular shapes. The overall scene is a dense collection of celestial objects.

A philosophical question:
What is the Fate of the Universe?


The Fate of the Universe can be determined from its history:



...And supernovae can be used as **standard candles** for this measurement



Expansion History of the Universe



Astrophysical Journal 1938

THE ABSOLUTE PHOTOGRAPHIC MAGNITUDE
OF SUPERNOVAE*

W. BAADE

ABSTRACT

A compilation of the photometric data for the 18 supernovae known at the end of 1937 is given. Former estimates have been replaced by photometric magnitudes after a redetermination of the magnitudes of comparison stars on the international system. The mean absolute photographic magnitude of the supernovae, derived from this material, is $M_{\max} = -14.3 \pm 0.42$ (m.e.) with a dispersion $\sigma_{M_{\max}} \approx 1.1$ mag. This result, together with the spectroscopic evidence, fully confirms the view that two classes of novae, common novae and supernovae, exist. Attention is drawn to the curious fact that 72 per cent of the known supernovae appeared in late-type spirals. B Cassiopeiae and the Crab nebula, which may have been galactic supernovae, are discussed.


When, after Ritchey's discovery of a nova in the spiral nebula NGC 6946 in 1917, novae were discovered in rapid succession in other extragalactic systems, a new way had been opened to settle the old question as to the constitution and the distances of these systems. The occurrence of novae in them afforded strong evidence for their stellar constitution. Moreover, their distances could be measured as soon as reliable values for the luminosities of the novae of our own galaxy were available. Nevertheless, the first applications of this method by H. D. Curtis and J. H. Oort were very satisfactory. Since the late 1920s, the data have been very

But supernovae were not quite good enough
“standard candles”



mid - 1980' s: Two new developments

1 “Type Ia” supernovae: a more standard candle




Panagia (1985)

Uomoto & Kirshner (1985)

Wheeler & Levreault (1985)

Two new

“Type Ia” supernovae
a more standard candle




Mid - 1980' s:

Two new developments

- 2 CCD detectors
& **computers** fast enough for image analysis


2

CCD detectors
& **computers** fast enough for image analysis




SN + Galaxy

—



Galaxy


=



SN

Luis Alvarez suggests
to Rich Muller that
it is time to re-do
Stirling Colgate's
robotic SN search

R. Muller:
Berkeley Automated
Supernova Search
*with C. Pennypacker
and S.P.*



MAY 8, 1986



MAY 17, 1986

SUPERNOVA IN M99 GALAXY IN VIRGO CLUSTER

Photo by : LBL SUPERNOVA SEARCH TEAM.

Why is the supernova measurement *not* easy?

1. Can they be found **far enough** -- and **enough of them** -- for cosmology?
Can they be found **early enough** to measure brightness over peak?
2. Can they be identified as Type Ia with spectra, despite how faint they will be?
Can their brightness be compared with nearby ones, despite greatly "redshifted" spectra?
3. Are the supernovae standard enough?
And how can one eliminate possible dust from diminishing their brightness?
4. Couldn't the supernovae evolve over 5 billion years?




Problems

with Type Ia Su

Rare

Random

Rapid




NB: Norgaard-Nielsen et al. searched intensively for over 2 years, and found just 1 Type Ia supernova several weeks past its peak.

can't schedule observations or plan discoveries at new moon

difficult to catch on the rise

Pennypacker & Perlmutter 1987 proposal:


A novel F/1 wide-field CCD camera
for the Anglo-Australian 4-m telescope (AAT)




...A big enough
telescope with a
wide enough field
to search for
 $z > 0.3$ Type Ia
supernovae in
100s of galaxies
with each image.


f855 7 01

Pennypacker & Perlmutter 1988 wide-field CCD camera at AAT





F855 7 01



2950.00

5555.46

3070.00

5755.46

F855 7 01



2050.00

5755
3070.00

January 1990 — November 1989

F855 7 1990:01:23



-50.0000 150.000

Problems

with Type Ia Supernovae as a tool for cosmology

Rare

$\sim 1 / 500$ years / galaxy



Random

can't schedule telescope time
or plan discoveries at new moon



Rapid

difficult to catch on the rise

Hamuy et al. (*Astronomical Journal* 1993),
describing the Calan/Tololo Search for supernovae at much lower redshifts:

“Unfortunately, the appearance of a SN is **not predictable**. As a consequence of this we **cannot schedule** the followup observations a priori, and we generally have to rely on **someone else’s telescope time**. This makes the execution of this project **somewhat difficult**.”



Random

can't schedule telescope time
or plan discoveries at new moon



Rapid

difficult to catch on the rise

Search Strategy

Perlmutter *et al* (1994)



Central Bureau for Astronomical Telegrams

INTERNATIONAL ASTRONOMICAL UNION

Postal Address: Central Bureau for Astronomical Telegrams

Smithsonian Astrophysical Observatory, Cambridge, MA 02138, U.S.A.

IAUSUBS@CFA.HARVARD.EDU or FAX 617-495-7231 (subscriptions)

BMARSDEN@CFA.HARVARD.EDU or DGREEN@CFA.HARVARD.EDU (science)

Phone 617-495-7244/7440/7444 (for emergency use only)

SUPERNOVAE

The Supernova Cosmology Project [S. Perlmutter, S. Deustua, G. Goldhaber, D. Groom, I. Hook, A. Kim, M. Kim, J. Lee, J. Melbourne, C. Pennypacker, and I. Small, Lawrence Berkeley Lab. and the Center for Particle Astrophysics; A. Goobar, Univ. of Stockholm; R. Pain, CNRS, Paris; R. Ellis and R. McMahon, Inst. of Astronomy, Cambridge; and B. Boyle, P. Bunclark, D. Carter, and M. Irwin, Royal Greenwich Obs.; with A. V. Filippenko and A. Barth (Univ. of California, Berkeley) at the Keck telescope; W. Couch (Univ. of N.S.W.) and M. Dopita and J. Mould (Mt. Stromlo and Siding Spring Obs.) at the Siding Spring 2.3-m telescope; H. Newberg (Fermi National Accelerator Lab.) and D. York (Univ. of Chicago) at the ARC telescope] report eleven supernovae found with the Cerro Tololo (CTIO) 4-m telescope in their 1995 High Redshift Supernova Search:

SN	1995 UT	R.A. (2000)	Decl.	R	Offset
1995aq	Nov. 19	0 29 04.26	+ 7 51 20.0	22.4	0".6 W, 1".4 S
1995ar	Nov. 19	1 01 20.41	+ 4 18 33.8	23.1	2".9 W, 0".5 S
1995as	Nov. 19	1 01 35.30	+ 4 26 14.8	23.3	0".7 W, 0".7 N
1995at	Nov. 20	1 04 50.94	+ 4 33 53.0	22.7	0".3 W, 0".4 S
1995au	Oct. 29	1 18 32.60	+ 7 54 03.5	20.7	1".4 E, 3".3 N
1995av	Nov. 20	2 01 36.75	+ 3 38 55.2	20.1	0".2 W, 0".0 N
1995aw	Nov. 19	2 24 55.54	+ 0 53 07.5	22.5	0".2 W, 0".2 S
1995ax	Nov. 19	2 26 25.80	+ 0 48 44.2	22.6	0".3 W, 0".2 S
1995ay	Nov. 20	3 01 07.52	+ 0 21 19.4	22.7	0".9 W, 1".4 S
1995az	Nov. 20	4 40 33.59	- 5 30 03.6	24.0	1".6 W, 1".7 N
1995ba	Nov. 20	8 19 06.46	+ 7 43 21.2	22.6	0".1 E, 0".2 N


The spectra (Keck, Nov. 26-28) are consistent with type-I supernovae (except SN 1995av, a probable type II) at the redshift of the host galaxy: $z = 0.45, 0.46, 0.49$ (preliminary type-I identification), $0.65, 0.16, 0.30, 0.4$ (supernova redshift only), $0.61, 0.48, 0.45, 0.39$. Photometry obtained on Nov. 21-23 at CTIO (A. Walker) and Nov. 23-27 at WIYN (D. Harmer, D. Willmarth) indicates that SNe

Why is the supernova measurement *not* easy?


1. Can they be found **far enough** -- and **enough of them** -- for cosmology?
Can they be found **early enough** to measure brightness over peak?
2. Can they be identified as Type Ia with spectra, despite how faint they will be?
Can their brightness be compared with nearby ones, despite greatly "redshifted" spectra?
3. Are the supernovae standard enough?
And how can one eliminate possible dust from diminishing their brightness?
4. Couldn't the supernovae evolve over 5 billion years?






And, in fact, the spectra do look like noise
...until you know what you are looking for.




Observed Wavelength (Å)










Type Ia
spectrum
redshifted
to $z = 0.42$







Type Ia
spectrum
redshifted
to $z = 0.5$






Type Ia
spectrum
redshifted
to $z = 0.55$







Type Ia
spectrum
redshifted
to $z = 0.55$





Standard
K corrections

Leibundgut, A&A (1990)



“Cross-Filter”
K corrections

Kim, Goobar, & Perlmutter (1995)




Why is the supernova measurement *not* easy?

1. Can they be found **far enough** -- and **enough of them** -- for cosmology?
Can they be found **early enough** to measure brightness over peak?
2. If found, they won't be bright enough to identify as Type Ia with spectrum.
And how can their brightness -- greatly redshifted -- be compared with nearby ones?
3. Are the supernovae standard enough?
And how can one eliminate possible dust from diminishing their brightness?
4. Couldn't the supernovae evolve over 5 billion years?




“Type Ia”?


Panagia (1985)




Leibundgut & Tammann
A&A (1990)



Nearby Type Ia Supernova



Nearby Type Ia Supernova



Observed dispersion of nearby
Type Ia peak brightness:

Branch & Miller (1993)

Vaughan, Branch, Miller, & S.P. (1995)

40% -- 50% **observed** dispersion
reduced to 30% dispersion by
selection based on color

Hamuy, Maza, Phillips, Suntzeff et al (1993)

“Calan/Tololo Supernova Search”

A beautiful, well-measured set of nearby supernova

now observed dispersion goes down to
~18% after color selection

Lightcurve Width-Luminosity Relation


Phillips:
(1993-)



CHARACTERIZED BY:

Decline Rate

Riess, Press, & Kirshner:
(1995-)



Light Curve Shape (LCS)



Perlmutter et al.:
(1996-)



Timescale "stretch factor"


$S > 1$: Broader / Slower
light curves are Brighter

$S < 1$: Narrower / Faster
light curves are Fainter



more redshift ↑


Type Ia Supernovae



Throw out *red* supernovae:
they are dimmed by dust
or peculiar

fainter →

Compare color distributions,
or correct each SN
individually for its
color, assuming a
dust color law.




Why is the supernova measurement *not* easy?

1. Can they be found **far enough** -- and **enough of them** -- for cosmology?
Can they be found **early enough** to measure brightness over peak?
2. Can they be identified as Type Ia with spectra, despite how faint they will be?
Can their brightness be compared with nearby ones, despite greatly "redshifted" spectra?
3. Are the supernovae standard enough?
And how can one eliminate possible dust from diminishing their brightness?
4. Couldn't the supernovae evolve over 5 billion years?



SN Ia Host Galaxies: Morphological Classification with HST/STIS Imaging



Why is the supernova measurement *not* easy?

1. Can they be found **far enough** -- and **enough of them** -- for cosmology?
Can they be found **early enough** to measure brightness over peak?
2. Can they be identified as Type Ia with spectra, despite how faint they will be?
Can their brightness be compared with nearby ones, despite greatly "redshifted" spectra?
3. Are the supernovae standard enough?
And how can one eliminate possible dust from diminishing their brightness?
4. Couldn't the supernovae evolve over five billion years?




Why is the supernova measurement *not* easy?

a “new”
problem


5. What if Einstein’s “Cosmological Constant” (Λ) exists? It will fight against gravity due to mass (M) in the universe

-- how can you tell if there is less M or more Λ or vice versa?





cosmological
constant
density




mass density

Lunar Calendar



50-100
Fields

Almost 1000
Galaxies per
Field




RESULT: ~12 Type Ia supernovae
discovered while still brightening,
at new moon




Scheduled Follow-Up
Spectroscopy at Keck



Scheduled Follow-Up
Imaging at Hubble,
Cerro Tololo,
WIYN, Isaac Newton



Hubble



Berkeley
Lab


WIYN


Isaac Newton


Keck

Cerro Tololo
with Tony Tyson & Gary
Bernstein's
"Big Throughput Camera"


redshift distribution of Type Ia supernovae as of 1998









Expansion History of the Universe




Expansion History of the Universe




Expansion History of the Universe



Expansion History of the Universe






...or a modification of Einstein's
Theory of General Relativity?


004-6961/2003/75(2)559\$4.95/35.00	559	2003 The American Phys
------------------------------------	-----	------------------------

Everybody talks about the dark energy,
but nobody does anything about it.


Expansion History of the Universe




Expansion History of the Universe



Union2 Compilation




Expansion History of the Universe







Carl
Pennypacker



Rich
Muller



Saul
(with hair)

Carl
Pennypacker

Heidi
Newberg

Warrick
Couch

Shane
Burns



Rich
Muller


Brian
Boyle




Heidi
Newberg









Warrick
Couch



NOVEMBER
1989




JANUARY
1990



Reynald
Pain



Alex
Filippenko



Brad
Schaefer



Pilar
Ruiz-Lapuente



Sebastian
Fabbro



Matthew
Kim




Ivan
Small



Isobel
Hook



Don
Groom



Susana
Deustua



Chris
Lidman



Circular
Central Bureau for Astronomical Telegrams
INTERNATIONAL ASTRONOMICAL UNION
Postal Address: Central Bureau for Astronomical Telegrams
Smithsonian Astrophysical Observatory, Cambridge, MA 0213
IAUSUBS@CFA.HARVARD.EDU or FAX 617-495-7231 (subscription
BMAUSDEN@CFA.HARVARD.EDU or DGREEN@CFA.HARVARD.EDU (scien
Phone 617-495-7244/7440/7444 (for emergency use only)

SUPERNOVAE
The Supernova Cosmology Project [S. Perlmutter, S. D.
Goldhaber, D. Groom, I. Hook, A. Kim, M. Kim, J. Lee, J.
C. Pennypacker, and I. Small, Lawrence Berkeley Lab. and
for Particle Astrophysics; A. Goobar, Univ. of Stockholm;
CNRS, Paris; R. Ellis and R. McMahon, Inst. of Astronomy
Cambridge; and B. Boyle, P. Bunclark, D. Carter, and M. I.
Royal Greenwich Obs.; with A. V. Filippenko and A. Barth (Univ. of
California, Berkeley) at the Keck telescope; W. Couch (Univ. of
N.S.W.) and M. Dopita and J. Mould (Mt. Stromlo and Siding Spring
Obs.) at the Siding Spring 2.3-m telescope; H. Newberg (Fermi
National Accelerator Lab.) and D. York (Univ. of Chicago) at the
ARC telescope] report eleven supernovae found with the Cerro Tololo
(CTIO) 4-m telescope in their 1995 High Redshift Supernova Search:

SN 1995 WT P.A. (2000) Decl. P. Offset






Andy
Fruchter



Nino
Panagia





Rob
Knop




Peter
Nugent



Greg
Aldering






Rob
Knop



Peter
Nugent



Greg
Aldering



Julia
Lee



Nelson
Nunes



Patricia
Castro



Robert
Quimby



David
Branch



Mario
Hamuy



José
Maza



Craig
Wheeler



Gustav
Tammann



